

Cover crop differentially affects arthropods, but not diseases, occurring on grape leaves in vineyards

F. VOGELWEITH^{1,2}  and D. THIÉRY²

¹ Abteilung Evolutionsbiologie, Johannes Gutenberg-Universität Mainz, Institut für Zoologie, 55128 Mainz, Germany; ² Institut National de la Recherche Agronomique (INRA), Unité Mixte de Recherche (UMR) 1065 Santé et Agroécologie du Vignoble (SAVE) Institut des Sciences de la Vigne et du Vin (ISVV), F-33883 Villenave d'Ornon Cedex, France
Corresponding author: Dr Fanny Vogelweith, email fanny.vogelweith@gmail.com

Abstract

Background and Aims: Cover crop has become a common management practice in viticulture. It improves the structure and protects the soil, enhances natural enemy populations and also reduces the vigour of the vine. Here, we investigated the effect of cover crop in vineyards on grapevine diseases and arthropods present on leaves.

Methods and Results: We measured the presence of the pathogens *Plasmopara viticola*, *Uncinula necator* and *Guignardia bidwellii* and monitored six common beneficial/pest arthropods: *Panonychus ulmi*, *Orthotydeus lambi*, *Typhlodromus pyri*, *Scaphoideus titanus* and *Phalangium opilio* in vineyards with cover crop or bare soil in 2014 and 2015. The density of the two pests *P. ulmi* and *S. titanus* and the beneficial mite *O. lambi* was lower in cover crop, while the density of the beneficial predatory mite *T. pyri* was higher in cover crop. We found no influence of cover crop on *P. opilio* and on the presence of grapevine diseases.

Conclusions: These findings point to a simple way to increase the population of beneficial arthropods and reduce pest arthropods, which could help integrated pest management programs.

Significance of the Study: To our knowledge, this is the first study monitoring the cover crop influence on both diseases and arthropods. It would help growers to optimise space management between rows to enhance their natural enemy populations while reducing pesticide application.

Keywords: black-rot, ground cover, leafhopper, mildew, mites

Introduction

Growers have used cover crop in orchards and vineyards for centuries (Ingels 1998). Abandoned in the 20th century for synthetic fertilisers, cover crop experienced a widespread resurgence in the 21st century, as a result of the growing interest in sustainable agriculture and biological control (Ingels 1998). Cover crop has been shown to slow down erosion, improve soil structure, increase the development of beneficial microorganisms, smother weeds and regulate minerals, moisture and water supply (Landis et al. 2000, Gurr et al. 2004, Silvestre et al. 2012, Irvin et al. 2016). Despite these benefits, cover crop can also have a detrimental effect on vine growth, yield and grape composition (Caspari et al. 1997, Silvestre et al. 2012).

Cover crop can enhance populations of beneficial insects in agricultural systems and improve pest control as a result (Landis et al. 2000, Gurr et al. 2004, Irvin et al. 2016). In vineyards, it often increases the population of natural enemies of pests, thus reducing spider mite and some leafhopper populations on grapes (Hanna et al. 1996, Costello and Daane 1998, English-Loeb et al. 2003). For instance, Costello and Daane (1998) demonstrated, over 4 years, a lower density of leafhopper nymphs in cover crop vineyards compared with that of control vineyards. In their study investigating both pests (e.g. thrips, leafhoppers and mites) and beneficial arthropods (e.g. earwigs and spiders), Irvin et al. (2016) found that cover crop caused an increase of generalist predators and a migration of beneficial arthropods as well as pests into the grape canopy. Ultimately, both the

efficiency and impact of cover crop in vineyards appear highly variable, thus requiring further study (Danne et al. 2010, Pérez-Alvarez et al. 2015).

Cover crop also creates microclimatic changes including a local decrease in temperature and an increase in humidity (Landis et al. 2000). Because temperature and humidity are key conditions for fungal growth, cover crop should also affect grape diseases. Downy and powdery mildew and, to a lesser extent, black-rot are three important leaf diseases causing considerable damage in vineyards (Ficke et al. 2002, Hoffman et al. 2002, Gadoury et al. 2003, Kennelly et al. 2005). These three fungi affect vine growth, yield and fruit composition leading to substantial losses and are managed principally with fungicides (Ficke et al. 2002, Hoffman et al. 2002, Gadoury et al. 2003, Kennelly et al. 2005). To our knowledge, no study has clearly tested the effect of cover crop on grapevine leaf diseases. This is surprising because cover crops are used with increased frequency in vineyards (Ingels 1998), and these diseases can be devastating (Valdés-Gómez et al. 2011), inducing the use of large amounts of fungicides [80% of pesticides used in French viticulture are motivated by grape downy and powdery mildews (Delière et al. 2015)]. By limiting the nitrogen available in the vine and thus reducing its vigour, cover crops could thus decrease to some extent both grape powdery and downy mildew (Hoffland et al. 2000, Mitchell et al. 2003). They could also increase, however, the moisture retention by leaves promoting the plant fungal diseases (Dake 1995, Singh et al. 2011).

In this study, we investigated the effect of cover crop on grapevine leaf diseases and arthropods living on leaves. We monitored three of the most common grapevine diseases in French vineyards and six species of arthropods considered as pests or beneficial organisms over 4 months in 2014 and 2015. Because of microclimatic changes induced by cover crop, we predicted a higher frequency of grape diseases as a result of cover crop management. Moreover, because cover crop increases the population of natural enemies, we predicted a higher density of beneficial arthropods and a lower density of pest arthropods in cover crop management.

Material and methods

Vineyard plot and disease management

The experiment took place in the experimental vineyard of INRA Research Center la Grande Ferrade (Villenave d'Ornon, France). The vineyard plot is planted to Merlot vines (clone 181) and was planted in 1991. The management of the vineyard is a double Guyot training system with leaf and top removal every 14 days, if necessary. The area of the plot is 1383 m² containing 16 rows of 42 vines (total of 672 vines) and a density of about 5700 vines/ha. The distance between rows is 1.60 m and between vines 1.10 m. In this experiment, we used only rows 2, 3 and 4. These vines were non-treated (for more than 10 years) against diseases or arthropod pests. In each of the three rows tested, we did not use the two outermost vines because of potential effects from the surrounding grape plots. The 40 vines used in each row alternate five vines with natural cover crop and five vines without cover crop (called bare soil).

Experimental design and sampling

We assessed the effect of cover crop on the presence of disease symptoms and on the presence and density of six species of arthropods present on the leaves. Data were collected on seven occasions from May to August in both 2014 and 2015 (a total of 14 replicates).

Because both diseases and arthropods can be affected by the different environmental conditions between years, we recorded precipitation, temperature and humidity from May to the end of August. In 2014, a monthly average of 2.10 mm of rain fell in the vineyard, the mean temperature per month was 19.14°C (minimum 14.48°C; maximum 24.48°C) and the humidity per month was 72.47% (minimum 48.33%; maximum 93.69%). In 2015, a monthly average of 0.65 mm of rain fell in the vineyard, the mean temperature per month was 21.60°C (minimum 14.89°C; maximum 27.67°C) and the humidity per month was 65.64% (minimum 38.67%; maximum 92.62%).

Grapevine arthropods on leaves. Three mite species (*Orthotydeus lambi*, *Panonychus ulmi* and *Typhlodromus pyri*), the leafhopper *Scaphoideus titanus* and the harvestman *Phalangium opilio* were monitored. These species are common in French vineyards and can be described as pests and/or beneficial arthropods (Sentenac 2011).

Phalangium opilio (Opiliones: Phalangidae) is the most common harvestman species in disturbed habitats, such as crops (Anderson 2012). This species is considered as an effective generalist predator in vineyards, controlling pest populations (Anderson 2012). Here, the number of harvestmen was visually counted on each vine, and 20 vines were assessed per treatment at each data collection. *Scaphoideus titanus* (Hemiptera: Cicadellidae) is a specialist leafhopper of

Vitis in Europe and is the only known vector of flavescence dorée, one of the most destructive diseases in European vineyards (Chuche and Thiéry 2014). Flavescence dorée is a wall-less intracellular bacterium restricted to phloem sieve tubes (Papura et al. 2012), highly pathogenic to several major grapevine cultivars and often lethal to the vine. Thus, *S. titanus* is considered a major pest in European vineyards (Chuche and Thiéry 2014). The number of leafhoppers was visually counted on five leaves chosen at random per vine, directly in the vineyard. Twenty vines per replicate were assessed.

After these measurements, five leaves per vine (total of 100 leaves sampled for each treatment per replicate) were haphazardly sampled and brought to the laboratory for mite species determination (Collyer 1982, Bolland et al. 1998, Zhang et al. 2001). Each leaf was brushed with a mite brush, and the mites were collected, counted and identified with a binocular microscope (magnification 20×). We obtained an average number of each species per vine. We identified and focused on three major mite species during this monitoring: *O. lambi*, *P. ulmi* and *T. pyri*. *Orthotydeus lambi* is a mycophagous mite and highly abundant in vineyards (Melidossian et al. 2005, English-Loeb et al. 2007). This species can be considered as beneficial in vineyards because *O. lambi* can reduce powdery mildew on foliage and fruit (English-Loeb et al. 2007). Contrary to *O. lambi*, *P. ulmi* is a phytophagous mite and considered as a pest in vineyards and orchards (Alston and Reding 2011). Populations of *P. ulmi*, however, can be controlled by the predatory mite *T. pyri* commonly used in biological control in vineyards (Marshall and Lester 2001). *Typhlodromus pyri* is also an interesting species because it can use the grape powdery mildew as an alternative food source (Pozzebon and Duso 2008).

For each vine, we counted the number of primary and secondary branches and the number of leaves on these branches. Each sampled leaf was transversely measured with a ruler. We obtained the number of leaves on each vine and the average size of the leaves (precision ± 0.01 mm). Thus, we corrected the number of leafhoppers and mites by both the number and the size of the leaves to obtain a density in cm². The vine phenology was recorded at each replicate following the BBCH-scale for grapevine (Lancashire et al. 1991).

Grapevine diseases on leaves. We studied the three more common grapevine diseases present on leaves in French vineyards: the grape downy mildew (*Plasmopara viticola*), the grape powdery mildew (*Ucinula necator*) and the black-rot (*Guignardia bidwellii*).

The grape downy mildew is caused by the oomycete *P. viticola* (Kennelly et al. 2005) and powdery mildew by the ascomycete *U. necator* (Gadoury et al. 2003). These two fungi require specific microclimates for infection (mild temperature with high humidity) (Gadoury et al. 2003, Kennelly et al. 2005). Once infected, the vine suffers numerous injuries, such as a reduction of the photosynthetic capacity caused by a premature defoliation (Pool et al. 1984) and a delay in berry development (Wan et al. 2007). Symptoms of *U. necator* appear as irregular chlorosis of grey-white with white powder on the leaf surface and as black net lines with white powder on the surface of the berry, stalk and tendril, and symptoms of *P. viticola* appear as yellowish, oily lesions on the leaf surface (Wan et al. 2007). The ascomycete *G. bidwellii* causes black-rot. The infection period of this fungus is determined by the correlations between leaf wetness duration and temperature (Jermini and Gessler 1996). Symptoms of *G. bidwellii* appear as spots on the

leaves, tan to reddish-brown circular lesions (Hoffman et al. 2002).

We estimated the presence of each disease by recording at random the symptoms on 15 leaves per vine. Twenty vines per treatment were assessed. An average presence of each disease on each vine was then calculated.

Statistical analysis. The software R v3.1.2 loaded with the packages *car*, *lme4* and *MASS* was used for all statistical analyses (R Development Core Team, 2011). The corrected number of arthropods on the leaves (harvestmen, leafhoppers and mites) was analysed using linear mixed models (*lmer*), in which treatments (bare soil or cover crop) and the years (2014 or 2015) were entered as explanatory categorical factors, while the vine number and the grapevine phenology were entered as a random effect.

The presence of each disease (*P. viticola*, *U. necator* and *G. bidwellii*) was tested using a generalised linear mixed model (*glmer*, with binomial distribution). In these models, the

treatments and the years were entered as explanatory categorical factors, while the vine number and the grapevine phenology were entered as a random effect.

All models first included interactions between the two explanatory factors and were then simplified stepwise by removing the non-significant interaction terms (all *P*-values > 0.08). Note that some non-significant interactions are presented here to allow direct comparisons between models, but their removal did not qualitatively change the results.

Results

Grapevine arthropods on leaves

The presence of cover crop in the vineyard shaped the density of the monitored mites, *P. ulmi*, *O. lambi* and *T. pyri*, and of the leafhopper *S. titanus* but not that of the harvestman *P. opilio* (Table 1, Figure 1). Specifically, the two pests *P. ulmi* and *S. titanus* were more dense in bare soil compared with that with

Table 1. Effect of the treatment (bare soil or cover crop) and the sampling year (2014 or 2015) on the density of leaf arthropods tested.

	<i>Panonychus ulmi</i>		<i>Orthotydeus lambi</i>		<i>Typhlodromus pyri</i>		<i>Scaphoideus titanus</i>		<i>Phalangium opilio</i>	
	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value	<i>F</i>	<i>P</i> -value
Treatment	9.53	0.002	17.13	<0.0001	5.17	0.023	15.82	<0.0001	0.26	0.612
Year	0.67	0.411	19.08	<0.0001	0.43	0.513	0.01	0.915	0.22	0.638
Treatment*year	0.02	0.894	6.29	0.012	0.44	0.506	0.28	0.594	1.16	0.281

Significant *P* values are in bold.




Figure 1. Effect of cover crop (■) on the number of arthropods on grapevine leaves compared with that of bare soil (□): (a) *Panonychus ulmi* (\pm SE); (b) *Orthotydeus lambi* (\pm SE) depending on the sampling year; (c) *Typhlodromus pyri* (\pm SE) and *Scaphoideus titanus* (\pm SE). The number in the bars represents the sample size. ***, *P* < 0.0001; *, *P* < 0.05; and n.s., non-significant.

Table 2. Effect of the treatment (bare soil or cover crop) and the sampling year (2014 or 2015) on the proportion of diseases on grapevine leaves.

	<i>Plasmopara viticola</i>		<i>Uncinula necator</i>		<i>Guignardia bidwellii</i>	
	F	P-value	F	P-value	F	P-value
Treatment	2.44	0.118	0.05	0.826	0.77	0.378
Year	55.52	<0.0001	16.29	<0.0001	136.14	<0.0001
Treatment*year	0.57	0.448	0.01	0.906	2.58	0.108

Significant *P* values are in bold.




Figure 2. Effect of cover crop (■) and on the proportion of grapevine diseases on leaves/vine compared with that of bare soil (□) for the years 2014 and 2015: (a) grape downy mildew (*Plasmopara viticola*; ±SE); (b) grape powdery mildew (*Uncinula necator*; ±SE); and (c) black-rot (*Guignardia bidwellii*; ±SE). The number in the bars represents the sample size. ***, *P* < 0.0001.

cover crop (Figure 1a,d), whereas the predatory mite *T. pyri* was more dense in cover crop compared with that in bare soil (Figure 1c). The density of *P. ulmi*, *T. pyri* and *S. titanus* was not affected by the sampling year (Table 1). The density of the beneficial mite *O. lambi*, however, was affected by the interaction between year and treatment (Table 1, Figure 1b): the density was higher in bare soil compared with that under cover crop in 2015, but not in 2014 (Figure 1b). Finally, the number of harvestmen on each vine was affected neither by the treatment (bare soil: mean ± SE = 0.28 ± 0.04 *P. opilio*/vine; cover crop: mean ± SE = 0.30 ± 0.04 *P. opilio*/vine) nor by the year (2014: mean ± SE = 0.15 ± 0.11 *P. opilio*/vine; 2015: mean ± SE = 0.18 ± 0.03 *P. opilio*/vine) (Table 1).

Grapevine diseases on leaves

The presence of cover crop in the vineyard did not affect the presence of the three grapevine diseases monitored, *P. viticola*, *U. necator* and *G. bidwellii* (Table 2). In contrast, all diseases were more frequent in 2015 than in 2014 (Table 2, Figure 2).

Discussion

The aim of this study was to determine the effect of cover crop on beneficial or pest arthropods and different grapevine diseases on leaves in a vineyard. Except for the harvestmen, the arthropods monitored were generally influenced by presence of cover crop but in different ways. *Panonychus ulmi*, *O. lambi* and *S. titanus* were more abundant in bare soil, and *T. pyri* was more abundant in cover crop. Grapevine diseases were not influenced by the presence of cover crop but were affected by the sampling year.

Cover crop reduced the density of the two pests *P. ulmi* and *S. titanus*, which is consistent with other studies on *P. ulmi* in orchards (Brown et al. 1997) or leafhoppers in vineyards (Costello and Daane 1998, 2003). Cover crop can indeed provide alternative food and shelter against natural enemies (Landis et al. 2000), shifting the density of arthropods from grapevine leaves to crop cover, as found in leafhoppers (Trivellone et al. 2012, 2013). Conversely, by favouring establishment of natural enemies, cover crop increases pests predation and then may further decrease arthropod numbers on grapevine leaves (Landis et al. 2000, Thomson and Hoffmann 2009, Tirello et al. 2013). Cover crop can also affect grapevine physiology (Caspari et al. 1997, Landis et al. 2000, Silvestre et al. 2012) and indirectly affect arthropod biology. For instance, Costello and Daane (2003) suggested that the lower density of leafhoppers in the cover crop treatment resulted from poorer host plant quality because of the competition between ground cover and grapevines.

Contrary to the other arthropod species, cover crop improved the density of the predatory mite *T. pyri*. In both

vineyards and orchards, cover crop promotes the colonisation of phytoseiid mites such as *T. pyri* (Duso et al. 2005, 2014, Aguilar-Fenollosa et al. 2011). Cover crop and adjacent vegetation around crop indeed increase the availability of alternative food sources, such as pollen (Duso et al. 2005, 2012). Because phytoseiidae mites, such as *T. pyri*, are known to be mobile (Tixier et al. 2000), it might be a good strategy for them to switch from vine leaves to adjacent vegetation when there is less food available. These results reinforced the positive involvement of cover crop on control of pests in vineyards. *Typhlodromus pyri* is an efficient biocontrol agent used in vineyards and orchards (Gerson et al. 2008, McMurtry et al. 2013). Cover crop appears to positively affect the population establishment of this predatory mite and negatively affects pests such as *P. ulmi* and *S. titanus* and thus favours biocontrol. Surprisingly, the density of the predator *P. opilio* was not influenced by cover crop. Contrary to spiders that are favoured by cover crop and display little mobility between vines (Costello and Daane 1998, 2003), we hypothesised that harvestmen can disperse more and then easily switch from one vine to the other.

The density of the mycophagous mite *O. lambi* was higher in 2015 compared with that in 2014. In 2015, the temperature over the season was higher than that in 2014 with a relatively high humidity level. These conditions should have favoured the development of fungi in the vineyard, not only on grapevine leaves but also in cover crop. Even though the variation of the density of tydeid mites between years is well-known (English-Loeb et al. 2007), tydeid mites, such as *O. lambi*, are also known to be abundant in the presence of grape downy and powdery mildew (Duso et al. 2005, Melidossian et al. 2005, English-Loeb et al. 2007), meaning that the density of *O. lambi* might be related to the presence of these diseases.

In 2015, the density of *O. lambi* was higher in bare soil compared with that in cover crop. As shown before for *P. ulmi* and *S. titanus*, cover crop can provide alternative food and shelter against natural enemies (Landis et al. 2000), inducing a shift of arthropod density from grapevine leaves to the cover crop.

The grape downy and powdery mildew and black-rot were not affected by the treatment but were influenced by the sampling year. Management practices such as vine training have been shown to affect diseases through microclimate changes. For example, Guyot training, in Bulgaria, favoured *B. cinerea* infection because of poor air circulation and temperature inversion, while the high leaf temperature of high-trained vines favoured powdery mildew infection (Draganov and Draganov 1976). Hence, we expected that microclimate changes induced by cover crop management affected grapevine leaf diseases. These conditions appear, however, to affect the grapevine diseases but at the year scale. In our experiment, the significant presence of diseases in 2015 might be due to the higher temperature compared with that in 2014 with a high level of humidity. In all three diseases monitored, the presence and intensity of these pathogens are indeed highly dependent on the year and consequently the climate (Spotts 1977, Jailloux et al. 1999, Madden et al. 2000, Harms et al. 2005). For instance, Jailloux et al. (1999) monitored grape powdery mildew during 5 years in a French vineyard of Merlot and found a strong influence of the April weather conditions (rainfall and temperature) on disease severity on berries by enabling good growth of the pathogen on leaves.

To conclude, we found that grapevine diseases on leaves were not affected by cover crop but by the climate between

years. Moreover, except for the harvestmen, leaf arthropods were differentially influenced by cover crop. These results could prove useful in vineyard management because we showed an increased density of the predatory mite *T. pyri* and a decreased density of pests such as *P. ulmi* and *S. titanus*. Such information would help growers to best manage the ground cover between rows to promote and enhance their natural enemy populations while reducing pesticide application.

Acknowledgements

We thank Delphine Binet, Lionel Delbac, Lionel Druelle and Pierre Sauris for their technical assistance. We also thank Romain Libbrecht and Joël Meunier for their insightful comments on the manuscript, Maximilian Körner for help with the English language and the anonymous reviewer for his helpful comments. This study was supported by CO-FREE European funding.

References

- Aguilar-Fenollosa, E., Ibáñez-Gual, M.V., Pascual-Ruiz, S., Hurtado, M. and Jacas, J.A. (2011) Effect of ground-cover management on spider mites and their phytoseiid natural enemies in clementine mandarin orchards (I): bottom-up regulation mechanisms. *Biological Control* **59**, 158–170.
- Alston, D.G. and Reding, M.E. (2011) European red mite. Utah pests website, fact sheet 1–4. <http://utahpests.usu.edu/ipm/html/fruits/fruit-insect-disease/euro-red-mites97> [Accessed 17/03/17].
- Anderson, M.S.R. (2012) The effect of landscape- and local-scale non-crop vegetation on arthropod pests and predators in vineyards. PhD Thesis. Lincoln University, Lincoln, New Zealand.
- Bolland, H., Gutierrez, J. and Flechtmann, C. (1998) World catalogue of the spider mite family (Acari: Tetranychidae) (Brill Academic Publishers: Leiden, The Netherlands).
- Brown, M.W., van der Zwer, T. and Glenn, D.M. (1997) Impact of ground cover plants on pest management in West Virginia, USA, apple orchards. *Horticultural Science* **24**, 39–44.
- Caspari, H.W., Neal, S. and Naylor, A. (1997) Cover crop management in vineyards to enhance deficit irrigation in a humid climate. *Acta Horticulturae* **449**, 131–320.
- Chuche, J. and Thiéry, D. (2014) Biology and ecology of the flavescence dorée vector *Scaphoideus titanus*: a review. *Agronomy for Sustainable Development* **34**, 381–403.
- Collyer, E. (1982) The phytoseiidae of New Zealand (Acarina) 1. The genera *Typhlodromus* and *Amblyseius* – keys and new species. *New Zealand Journal of Zoology* **9**, 185–206.
- Costello, M.J. and Daane, K.M. (1998) Effects of cover cropping on pest management: arthropods. Ingels, C.A., Bugg, R.L., McGourty, G.T. and Christensen, L.P., eds. *Cover cropping in vineyards: a grower's handbook* (University of California, Division of Agriculture and Natural Resources: Oakland, CA, USA) pp. 93–106.
- Costello, M.J. and Daane, K.M. (2003) Spider and leafhopper (*Erythronaura* spp.) response to vineyard ground cover. *Environmental Entomology* **32**, 1085–1098.
- Dake, G.N. (1995) Diseases of ginger (*Zingiber officinale* Rosc.) and their management. *Journal of Spices and Aromatic Crops* **4**, 40–48.
- Danne, A., Thomson, L.J., Sharley, D.J., Penfold, C.M. and Hoffmann, A.A. (2010) Effects of native grass cover crops on beneficial and pest invertebrates in Australian vineyards. *Environmental Entomology* **39**, 970–978.
- Delière, L., Cartolaro, P., Léger, B. and Naud, O. (2015) Field evaluation of an expertise-based formal decision system for fungicide management of grapevine downy and powdery mildews. *Pest Management Science* **71**, 1247–1257.
- Draganov, D. and Draganov, G. (1976) Influence of phytoclimate on several fungal diseases attacking high- and low-trained vines (translated from Bulgarian). *Gradinar Lozar Nauk* **13**, 96–102.
- Duso, C., Ahmad, S., Tirello, P., Pozzebon, A., Klaric, V., Baldessari, M., Malagnini, V. and Angeli, G. (2014) The impact of insecticides applied in apple orchards on the predatory mite *Kampimodromus aberrans* (Acari: Phytoseiidae). *Experimental and Applied Acarology* **62**, 391–414.
- Duso, C., Pozzebon, A., Capuzzo, C., Malagnini, V., Otto, S. and Borgo, M. (2005) Grape downy mildew spread and mite seasonal abundance

- in vineyards: effects on *Tydeus caudatus* and its predators. *Biological Control* **32**, 143–154.
- Duso, C., Pozzebon, A., Kreiter, S., Tixier, M.S. and Candolfi, M. (2012) Management of phytophagous mites in European vineyards. Bostanian, N.J., Vincent, C. and Isaacs, R., eds. *Arthropod management in vineyards: pests, approaches, and future directions* (Springer: Dordrecht, The Netherlands) pp. 191–217.
- English-Loeb, G., Rhainds, M., Martinson, T. and Ugine, T. (2003) Influence of flowering cover crops on *Anagrus parasitoids* (Hymenoptera : Mymaridae) and *Erythroneura* leafhoppers (Homoptera : Cicadellidae) in New York vineyards. *Agricultural and Forest Entomology* **5**, 173–181.
- English-Loeb, G., Norton, A.P., Gadoury, D., Seem, R. and Wilcox, W. (2007) Biological control of grape powdery mildew using mycophagous mites. *Plant Disease* **91**, 421–429.
- Ficke, A., Gadoury, D.M. and Seem, R.C. (2002) Ontogenic resistance and plant disease management: a case study of grape powdery mildew. *Phytopathology* **92**, 671–675.
- Gadoury, D.M., Seem, R.C., Ficke, A. and Wilcox, W.F. (2003) Ontogenic resistance to powdery mildew in grape berries. *Phytopathology* **93**, 547–555.
- Gerson, U., Smiley, R.L. and Ochoa, R. (2008) *Mites (Acari) for pest control* (Blackwell Science: Oxford, England).
- Gurr, G.M., Scarratt, S.L., Wratten, S.D., Berndt, L. and Irvin, N. (2004) Ecological engineering, habitat manipulation and pest management. Gurr, G.M., Wratten, S.D. and Altieri, M.A., eds. *Ecological engineering for pest management: advances in habitat manipulation for arthropods* (CSIRO Publishing: Collingwood, Vic., Australia) pp. 1–12.
- Hanna, B.R., Zalom, F.G. and Elmore, C.L. (1996) Integrating cover crops into vineyards. *Grape Grower* **27**, 26–43.
- Harms, M., Holz, G., Hoffmann, C., Lipps, H. and Silvanus, W. (2005) Occurrence of *Guignardia bidwellii*, the causal fungus of black rot on grapevine, in the vine growing areas of Rhineland-Palatinate, Germany. Proceedings international symposium on introduction and spread of invasive species; 9–11 June 2005; Berlin, Germany. http://p11631.typo3server.info/fileadmin/alte_Webseiten/Invasive_Symposium/articles/030_Harms.pdf
- Hoffland, E., Jeger, M.J. and van Beusichem, M.L. (2000) Effect of nitrogen supply rate on disease resistance in tomato depends on the pathogen. *Plant and Soil* **218**, 239–247.
- Hoffman, L.E., Wilcox, W.F., Gadoury, D.M. and Seem, R.C. (2002) Influence of grape berry age on susceptibility to *Guignardia bidwellii* and its incubation period length. *Phytopathology* **92**, 1068–1076.
- Ingels, C.A. (1998) *Cover cropping in vineyards: a grower's handbook* (University of California, Division of Agriculture and Natural Resources: Berkeley, CA, USA).
- Irvin, N.A., Bistline-East, A. and Hoddle, M.S. (2016) The effect of an irrigated buckwheat cover crop on grape vine productivity, and beneficial insect and grape pest abundance in southern California. *Biological Control* **93**, 72–83.
- Jailoux, F., Willocquet, L., Chapuis, L. and Froidefond, G. (1999) Effect of weather factors on the release of ascospores of *Uncinula necator*, the cause of grape powdery mildew, in the Bordeaux region. *Canadian Journal of Botany* **77**, 1044–1051.
- Jermi, M. and Gessler, C. (1996) Epidemiology and control of grape black rot in Southern Switzerland. *Plant Disease* **80**, 322–325.
- Kennelly, M.M., Gadoury, D.M., Wilcox, W.F., Magarey, P.A. and Seem, R.C. (2005) Seasonal development of ontogenic resistance to downy mildew in grape berries and rachises. *Phytopathology* **95**, 1445–1452.
- Lancashire, P.D., Bleiholder, H., Van Den Boom, T., Langelüddeke, P., Stauss, R., Weber, E. and Witzemberger, A. (1991) A uniform decimal code for growth stages of crops and weeds. *Annals of Applied Biology* **119**, 561–601.
- Landis, D.A., Wratten, S.D. and Gurr, G.M. (2000) Habitat management to conserve enemies of arthropod pests in agriculture. *Annual Review of Entomology* **45**, 175–201.
- Madden, L.V., Ellis, M.A., Lalancette, N., Hughes, G. and Wilson, L.L. (2000) Evaluation of a disease warning system for downy mildew of grapes. *Plant Disease* **84**, 549–554.
- Marshall, D. and Lester, P. (2001) The transfer of *Typhlodromus pyri* on grape leaves for biological control of *Panonychus ulmi* (Acari : Phytoseiidae, Tetranychidae) in vineyards in Ontario, Canada. *Biological Control* **20**, 228–235.
- McMurtry, J.A., Moraes, G.J.D.E. and Sourassou, N.F. (2013) Revision of the lifestyles of phytoseiid mites (Acari : Phytoseiidae) and implications for biological control strategies. *Systematic and Applied Acarology* **18**, 297–320.
- Melidossian, H.S., Seem, R.C., English-Loeb, G., Wilcox, W.F. and Gadoury, D.M. (2005) Suppression of grapevine powdery mildew by a mycophagous mite. *Plant Disease* **89**, 1331–1338.
- Mitchell, C.E., Reich, P.B., Tilman, D. and Groth, J.V. (2003) Effects of elevated CO₂, nitrogen deposition, and decreased species diversity on foliar fungal plant disease. *Global Change Biology* **9**, 438–451.
- Papura, D., Burban, C., van Helden, M., Giresse, X., Nusillard, B., Guillemaud, T. and Kerdelhué, C. (2012) Microsatellite and mitochondrial data provide evidence for a single major introduction for the nearctic leafhopper *Scaphoideus titanus* in Europe. *PloS One* **7**(5), e3688.
- Pérez-Alvarez, E.P., Garcia-Escudero, E. and Peregrina, F. (2015) Soil nutrient availability under cover crops: effects on vines, must, and wine in a Tempranillo vineyard. *American Journal of Enology and Viticulture* **66**, 311–320.
- Pool, R.M., Pearson, R.C., Welser, M.J., Lakso, A.N. and Seem, R.C. (1984) The influence of powdery mildew on the yield and growth of Rosette grapevines. *Plant Disease* **68**, 590–593.
- Pozzebon, A. and Duso, C. (2008) Grape downy mildew *Plasmopara viticola*, an alternative food for generalist predatory mites occurring in vineyards. *Biological Control* **45**, 441–449.
- R Development Core Team (2011), R: a language and environment for statistical computing. Vienna, Austria: the R Foundation for Statistical Computing. ISBN: 3–900051–07–0. Available online at <http://www.R-project.org/>.
- Sentenac, G. (2011) *La faune auxiliaire des vignobles de France* (France Agricole: Paris, France).
- Silvestre, J.C., Canas, S. and Brazao, J. (2012) Influence of timing and intensity of deficit irrigation on vine vigour, yield and berry and wine composition of “Tempranillo” in southern Portugal. *Acta Horticulturae* **931**, 193–201.
- Singh, R., Gangwar, S., Singh, R., Pandey, R. and Kalra, A. (2011) Medicinal plant *Coleus forskohlii* Briq.: disease and management. *Medicinal Plants* **3**, 1–7.
- Spotts, R.A. (1977) Effect of leaf wetness duration and temperature on the infectivity of *Guignardia bidwellii* on grape leaves. *Phytopathology* **77**, 1378.
- Thomson, L.J. and Hoffmann, A.A. (2009) Vegetation increases the abundance of natural enemies in vineyards. *Biological Control* **49**, 259–269.
- Tirello, P., Pozzebon, A. and Duso, C. (2013) The effect of insecticides on the non-target predatory mite *Kampimodromus aberrans*: laboratory studies. *Chemosphere* **93**, 1139–1144.
- Tixier, M.S., Kreiter, S., Croft, B.A. and Auger, P. (2000) Colonisation of vineyards by phytoseiid mites: their dispersal patterns in plot and their fate. *Experimental and Applied Acarology* **24**, 191–211.
- Trivellone, V., Paltrinieri, L.P., Jermi, M. and Moretti, M. (2012) Management pressure drives leafhopper communities in vineyards in Southern Switzerland. *Insect Conservation and Diversity* **5**, 75–85.
- Trivellone, V., Jermi, M., Linder, C., Cara, C., Delabays, N. and Baumgartner, J. (2013) Rôle de la flore du vignoble sur la distribution de *Scaphoideus titanus*. *Revue suisse Viticulture, Arboriculture, Horticulture* **45**, 222–228.
- Valdés-Gómez, H., Gary, C., Cartolaro, P., Lolas-Caneo, M. and Calonne, A. (2011) Powdery mildew development is positively influenced by grapevine vegetative growth induced by different soil management strategies. *Crop Protection* **30**, 1168–1177.
- Wan, Y., Schwaninger, H., He, P. and Wang, Y. (2007) Comparison of resistance to powdery mildew and downy mildew in Chinese wild grapes. *Vitis* **46**, 132–136.
- Zhang, Z.-Q., Bejakovich, D. and Martin, N.A. (2001) *Key to Tydeidae of New Zealand*. Ministry of Agriculture and Forestry Science Policy, Project FMA171. Landcare Research Contract Report: LC0001/118 (Landcare Research, Auckland, New Zealand) 36 pp.

Manuscript received: 07 November 2016

Revised manuscript received: 21 March 2017

Accepted: 03 April 2017